

AD-A263 396

AD_____

CONTRACT NO: DAMD17-88-C-8160

TITLE: UTILIZATION OF SIMIAN IMMUNODEFICIENCY VIRUS (SIV)

INFECTION FOR SUBHUMAN PRIMATES TO EVALUATE EXPERIMENTAL

CHEMOTHERAPY AND VACCINES

SUBTITLE: Determination of the ID50 of Frozen Stocks of Two

Strains of Simian Immunodeficiency Virus in Macaca

<u>Mulatta</u>

PRINCIPAL INVESTIGATOR: Dr. Preston Marx

CONTRACTING ORGANIZATION: The Regents of New Mexico

State University

Box 3699

Las Cruces, NM 88003-0001

REPORT DATE: July 1, 1992

TYPE OF REPORT: Final Report, Task Order II



PREPARED FOR: U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND

Fort Detrick, Frederick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;

distribution unlimited

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

93 4 20 050

REPORT DOCUMENTATION PAGE

Form Approved
OMB No 0704-0188

Public reporting burden for this callection of information is estimated to werage. Industrial per response including the time for reviewing instructions searching existing data sources gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for information Department and expects. 1215 Jefferson Davis Highway, Suite 1204, Arington, VA. 2202-4302, and to the Office of Management and Bugget, Page-mork Reduction Project (0704-0788). Washington, DC 20503.

| | 02-4302 and to the Office of Management and b | | | | |
|---|---|--|---|--|--|
| 1. AGENCY USE ONLY (Leave bla | · · | 3. REPORT TYPE AND D | | | |
| 4. TITLE AND SUBTITLE | 1 July 1992 | Final Report | , Task II FUNDING NUMBERS | | |
| Utilization of Sim Infection for Subh | ian Immunodeficiend uman Primates to Ev therapy and Vaccine | cy Virus (SIV) valuate es | | | |
| 7. PERFORMING ORGANIZATION N | AME(S) AND ADDRESS(ES) | 8. | PERFORMING ORGANIZATION | | |
| Box 3699 Las Cruces, New Me | | | REPORT NUMBER | | |
| | SENCY NAME(S) AND ADDRESS(ES) | | L SPONSORING / MONITORING AGENCY REPORT NUMBER | | |
| U.S. Army Medical Fort Detrick Frederick, Marylan | Research & Developm d 21702-5012 | ent Command | | | |
| 11 STIPPLEMENTARY NOTES | Subtitle: Determin | nation of the I | D50 of Frozen | | |
| Mulatta | ins of Simian Immu | | | | |
| 12a. DISTRIBUTION AVAILABILITY | * | 12 | b DISTRIBUTION CODE | | |
| Approved for publidistribution unlim | ited | | | | |
| 13. ABSTRACT (Maximum 200 work | ds) | | | | |
| The purpose of this research is to determine the infectivity of the two strains of Simian Immunodeficiency Virus (SIVmac 251 and SIVmac 239) in rhesus macaques. The data obtained from this study will provide information for the design of future lentiviral vaccine studies involving rhesus macaques and one of the above strains. | | | | | |
| 14. SUBJECT TERMS | | | 15 NUMBER OF PAGES | | |
| | nology; RA I; Task 'irus; Lab Animals; | | 16 PRICE CODE | | |
| 17. SECURITY CLASSIFICATION OF REPORT | 18 SECURITY CLASSIFICATION OF THIS PAGE | 19 SECURITY CLASSIFICAT OF ABSTRACT | ION 20. LIMITATION OF ABSTRACT | | |
| Unclassified | Unclassified | Unclassified | Unlimited | | |

FOREWORD

In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animal Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

Citations of commercial organizations and trade names in this report do not constitute an official Department of the Army endorsement or approval of the products or services of these organizations.

DITO GOLLENY INCPROTED 1

| Acces | sion For | ······································ |
|---------------|----------------------------|--|
| DTIC Unana | GRARI TAB counced fication | |
| By Distr | ibution/ | |
| Avai | lebility | |
| Dist | Avail and Special | |

CONTRACT NO. DAMD 17-88-C-8160 FINAL REPORT TASK ORDER II

Title:

Determination of the ID50 of Frozen Stocks of Two Strains of Simian Immunodeficiency Virus in Macaca mulatta.

Objective:

The purpose of this research is to determine the infectivity of the two strains of Simian Immunodeficiency Virus (SIVmac 251 and SIVmac 239) in rhesus macaques. The data obtained from this study will provide information for the design of future lentiviral vaccine studies involving rhesus macaques and one of the above strains.

Study Design:

On April 17, 1991, 38 candidate rhesus monkeys between the age of 18 months and 4 years were selected for the study. Two of these animals served as replacement if needed. These animals were observed daily and health observations were recorded during the entire baseline period, including body weights, body temperature, heart beats as well as clinical evaluations (CBC and SMAC). All 38 animals were tested negative for SRV (type D retrovirus). The virus inoculum was received from the Department of the Army in June, 1991. Serum from each animal was collected and frozen for serologic evaluation as baseline control before the inoculation on June 25, 1991.

Experimental Design:

1. Animal Inoculation:

Animals were inoculated by Dr. Rick Miller, D.V.M. on June 25, 1991. All animals received 1ml of the respective inoculum via the saphenous vein as designated in the table below.

| Viral Inoculum | Group | Log ₁₀ Dil. of Inoculum | No. of Anim. Per Group | Animals Designation |
|-------------------|-------|---------------------------------------|---------------------------------|------------------------|
| SIVmac 251 | 1 | 10 -2.5 | 3 | 5544, 5530, 88C |
| | 2 | 10 -3.5 | 4 | 5606, 92B, 5511, 5600 |
|] | 3 | 10 -4.5 | 4 | 81C, 5316, 3D, 105C |
| | 4 | 10 -5.5 | 4 | 5320, 5601, 110C, 5567 |
| | 5 | 10 -6.5 | 3 | 5568, 26C, 189B |
| SIVmac 239 | 1 | 10 -3.5 | 3 | 150B, 5561, 146B |
| | 2 | 10 -4.5 | 4 | 5549, 131B, 5593, 5543 |
| | 3 | 10 -5.5 | 4 | 101C, 5575, 5550, 5309 |
| | 4 | 10 -6.5 | 4 | 5319, 5605, 20D, 209B |
| | 5 | 10 -7.5 | 3 | 5603, 117B, 114C |

In addition to the cocultivation at PRL, a schedule of shipment of whole blood specimens from monkeys to be sent to sponsor/collaborators via overnight delivery was as follows:

| Date | Animal Group | PRL | Walter Reed Specimen |
|---------|-------------------|--------|----------------------|
| 6/25/91 | All animals, | Plasma | |
| Day 0 | groups 251 & 239 | | |
| 7/2/91 | All animals, | CBC | |
| Week 1 | groups 251 & 239 | | <u> </u> |
| 7/9/91 | All animals | Cocult | |
| Week 2 | groups 251 & 239 | | |
| 7/22/91 | 251 animals only | | 10ml heparinized |
| Week 4 | | | blood |
| 7/29/91 | 239 animals only | | 10ml heparinized |
| Week 5 | | | blood |
| 8/5/91 | 251 animals only, | | 10ml heparinized |
| Week 6 | groups 3, 4, 5 | | blood |
| 8/12/91 | 239 animals only, | | 10ml heparinized |
| Week 7 | groups 3, 4, 5 | | blood |
| 8/19/91 | 251 animals only, | | 10ml heparinized |
| Week 8 | groups 3, 4, 5 | | blood |
| 8/26/91 | 239 animals only, | | 10ml heparinized |
| Week 9 | groups 3, 4, 5 | | blood |

2. Virus Isolation:

Methods:

At day 14 post inoculation, virus isolation was performed. Peripheral blood mononuclear cells (PBMC) were separated from heparinized whole blood using lymphocyte separation medium (LSM) gradients. The PBMC were cocultivated with 2X106/ml CEMX174 (B cell and T cell hybrid) in complete RPMI 1640 medium supplemented with 10% Fetal Bovine Serum, 20mM L-glutamine, penicillin (100 μ g/ml)/streptomycin (100 μ g/ml). Cell free culture fluids were harvested once weekly and assayed for reverse transcriptase (RT) activity. All cultures were maintained for eight weeks or until positive for RT activity.

Results:The results are summarized in the table below:

| Virus Inoculum | Log 10 dilution | Animal Number | Isolation |
|----------------|-----------------|---------------|-----------|
| SIVmac 251 | 10 -2.5 | 5544 | + |
| | | 5530 | + |
| | | 88C | + |
| | 10 -3.5 | 92B | + |
| | | 5606 | + |
| | | 5511 | + |
| 1 | | 5600 | + |
| | 10 -4.5 | 81C | + |
| | | 5316 | - |
| | | 3D | - |
| | | 105C | + |
| | 10 -5.5 | 5320 | - |
| | | 5601 | + |
| | | 110C | - |
| | | 5567 | - |
| | 10 -6.5 | 5568 | - |
| | | 26C | - |
| | | 189B | - |
| SIVmac 239 | 10 -3.5 | 150B | + |
| | | 5561 | + |
| | | 146B | + |
| | 10 -4.5 | 5549 | + |
|] | | 131B | + |
| 1 | | 5593 | + |
| 1 | | 5543 | + |
| 1 | 10 -5.5 | 101C | - |
| | | 5575 | - |
| | | 5550 | + |
| | | 5309 | - |
| | 10 -6.5 | 5319 | - |
| | | 5605 | - |
| | | 20D | + |
| | | 209B | - |
| | 10 -7.5 | 5603 | - |
| | | 117B | - |
| | | 114C | - |

One week after cocultivation with CEMX174 cells, animals 131B and 5549 of the SIVmac 239 group ($10^{-4.5}$) were detected virus positive by RT assay. At week 3, animals 88C (SIVmac 251, $10^{-2.5}$), 5606 (251, $10^{-3.5}$) and 20D (239, $10^{-6.5}$) showed positive RT activity. By the fourth week, animals of the 251 group 5544, 5530 ($10^{-2.5}$), 5600 ($10^{-3.5}$), and 81C ($10^{-4.5}$) were detected RT positive. Animals 92B and 5511 ($10^{-3.5}$) of the 251 group and 150 B, 5561, 146B ($10^{-3.5}$) and 5550 ($10^{-5.5}$) of the 239 group turned positive at sixth and seventh week. During the final week of cocultivation, 105C ($10^{-4.5}$) and 5601 ($10^{-5.5}$) of the 251 group were virus positive. The rest of the animals remained virus negative by RT assay.

Animal 131B (239, 10 -4.5) was necropsied on July 14, 1991 by Dr. Lackner, PRL pathologist. At the time of necropsy, 10ml heparinized blood was collected for virus isolation. The PBMC were separated and cocultivated with CEMX174 cells for SIV. Two week post inoculation and at the time of death, this animal was determined virus positive by reverse transcriptase assay. Also at the time of necropsy, swabs of jejunum, stomach, colon and pericardium were cultured for Shigella and other pathogenic microorganisms. Only Shigella flexneri serogroup B was isolated in the jejunum.

TASK ORDER II-EXTENSION

In October, 1991, the Department of the Army requested to extend the Task Order II including challenge of the animals that were determined to be SIV negative after the first inoculation in June, 1991.

Before the virus challenge, all the study animals were bled for a final bleed (10ml and 5ml heparinized blood). The 10ml blood specimens were shipped to Dr. Lewis, Henry Jackson Foundation. The 5ml specimens were shipped to Dr. Ansari of Emory University Medical Center in Atlanta, Georgia, as requested by Dr. Lewis.

After the final bleed, inguinal lymph node biopsies were performed on the 251 monkeys: 5316, 105C, 3D, 5320, 5601, 5567, 5568, 26C, 81C, 189B and 110C, and on the 239 monkeys: 101C, 5575, 5550, 5309, 5319, 5605, 20D, 209B, 5603, 117B and 114C. The biopsy tissues were shipped to Dr. Lewis.

The SIV positive animals 88C, 92B, 5605, 81C, 5511 and 5600 were necropsied. The sterile tissues collected from axillary lymph nodes, inguinal lymph nodes, mesenteric lymph nodes, spleen and thymus. The remaining tissues were fixed in formalin. All tissues were sent to Dr. Lewis.

The animals 5544, 5530, 105C and 5601 of the SIVmac 251 group were held for six months to observe for disease outcome. These animals were bled at monthly intervals and the heparinized blood specimens were shipped to Henry Jackson Foundation. Additional peripheral blood and sera were collected for CBC and clinical analyses each time. The remaining confirmed SIV positive monkeys 150B, 5561, 146B, 5549, 5593, 5550, 20D, 101C and 5543 were returned to PRL for disposition.

The SIV-negative animals from the 251 and 239 groups were challenged on January 27, 1992. The animals included:

| SIVmac 251 | SIVmac 239 |
|------------|------------|
| 5316 | 5575 |
| 3D | 5309 |
| 5320 | 5319 |
| 110C | 5605 |
| 5567 | 209B |
| 5568 | 5603 |
| 26C | 117B |
| 189B | 114C |

The viral inoculum was diluted as follows:

SIVmac 251-4/19/91 dilutions:

| Dilution | 1:30 | 1:300 | 1:3000 |
|----------|--------|--------|--------|
| PBS | 2.9ml | 9.0ml | 9.0ml |
| Virus | *0.1ml | §1.0ml | 1.0ml |

SIVmac 239-10/1/90 dilutions:

| Dilution | 1:30 | 1:300 | 1:3000 | 1:30,000 |
|----------|--------|--------|---------------|----------|
| PBS | 2.9ml | 9.0ml | 9.0 ml | 9.0ml |
| Virus | *0.1ml | §1.0ml | §1.0ml | 1.0ml |

^{*}from virus obtained from R. Desrosiers

Each animal received 1ml of the viral inoculum via the left saphenous vein. The 251 animals received 1ml of the 1:3000 dilution each, while the 239 group animals received the 1:30,000 dilution each. Ten ml of heparinized blood was drawn from each animal at the time of challenge to be sent to Henry Jackson Foundation. Additional peripheral blood was collected for CBC and clinical chemistry. Following the virus challenge, all the animals were held for two months. Weekly blood draw of 10ml heparinized blood from each animal was sent to Henry Jackson Foundation.

The table below shows the schedule of the bleed:

| Week | Date | HJF 10ml hep. blood | CBC | SMAC | Physical | O&P |
|------|----------|------------------------|-----|------|----------|-----|
| 0 | *1/27/92 | X | X | X | X | X |
| 1 | 2/3/92 | X | | | | |
| 2 | 2/10/92 | X | X | l x | | |
| 3 | 2/18/92 | X | X | | X | |
| 4 | 2/24/92 | X | X | X | X | X |
| 5 | 3/2/92 | X | X | | X | |
| 6 | 3/5/92 | X | X | X | X | |
| 7 | 3 3/92 | X | X | | | |
| 8_ | , 23/92 | X | X | | X | |

^{*}day of challenge

Animals 5601 and 189B of the 251 group were necropsied before the end of the two month period due to declining health. Sterile tissues such as spleen, thymus, axillary, inguinal and mesenteric lymph nodes, heparinized blood were collected for

Sfrom previous dilution

Dr. Lewis. The remaining tissues were fixed in formalin and shipped to Dr. Lewis. A final CBC, clinical chemistry and blood cultures were also performed.

Four months after the challenge, the animals were scheduled to have axillary lymph node biopsies. All the challenged animals except 5568 underwent the procedure. Physicals and blood chemistry were done before the biopsy. Each biopsy tissue was divided for sterile specimen, frozen and paraffin blocks. Animal 5568 was necropsied due to declining health.

The remaining SIVmac 251 animals 3D, 26C, 110C, 5316, 5320, 5567, 5544, 5530 and 105C were requested by the Department of the Army to be released and transported to Henry Jackson Foundation upon completion of the study contract.

The remaining SIVmac 239 animals 56.33, 117B, 5575, 5309, 5319, 5605 209B and 114C were necropsied. Weights of pooled axillary lymph nodes, pooled inguinal lymph nodes, the ileocecal lymph node, and the spleen were recorded. Tissue samples of spleen, thymus, axillary lymph nodes, inguinal lymph nodes, mesenteric lymph nodes and the ileocecal valve were collected sterile. Small duplicate samples of the same tissues were processed for both frozen and paraffin blocks. A complete set of tissues including the CNS and GI tract was fixed in formalin and was shipped to Dr. Lewis with the rest of the sterile tissues.